

**2021**

**MATHEMATICS — HONOURS**

**Paper : CC-8**

**(Riemann Integration and Series of Functions)**

**Full Marks : 65**

*The figures in the margin indicate full marks.*

*Candidates are required to give their answers in their own words  
as far as practicable.*

$\mathbb{N}$ ,  $\mathbb{R}$  denote the sets of natural numbers and real numbers respectively.

1. Answer all the following multiple choice questions having only one correct option. Choose the correct option and justify : (1+1)×10
- (a) Let  $f : [a, b] \rightarrow \mathbb{R}$  be a bounded function and  $P, Q$  are partitions of  $[a, b]$  such that  $P$  is a refinement of  $Q$ . Then,
- |                              |                               |
|------------------------------|-------------------------------|
| (i) $L(P, f) \leq L(Q, f)$   | (ii) $L(P, f) \leq U(Q, f)$   |
| (iii) $U(P, f) \leq L(Q, f)$ | (iv) $U(P, f) \geq U(Q, f)$ . |
- (b) Let  $f : [0, 3] \rightarrow \mathbb{R}$  be defined by  $f(x) = [x]$ , where  $[x]$  denotes the greatest integer not exceeding  $x$ . Then,
- (i)  $f$  is not Riemann integrable on  $[0, 3]$ .
- (ii)  $f$  is Riemann integrable on  $[0, 3]$  and  $\int_0^3 f = 0$ .
- (iii)  $f$  is Riemann integrable on  $[0, 3]$  and  $\int_0^3 f = 2$ .
- (iv)  $f$  is Riemann integrable on  $[0, 3]$  and  $\int_0^3 f = 3$ .
- (c) Identify the incorrect statement :
- (i) Any subset of a negligible set is negligible.
- (ii) Any enumerable set of real numbers is negligible.
- (iii) Countable union of negligible sets is negligible.
- (iv) If the set of points of discontinuity of a real-valued function is negligible, then the function is monotonic.

**Please Turn Over**

(d) Let  $f : [0, 4] \rightarrow \mathbb{R}$  be defined by  $f(x) = x^4 - 4x^3 + 10$  and  $P = \{1, 2, 3, 4\}$ . Then,

(i)  $U(P, f) = -40$

(ii)  $L(P, f) = 11$

(iii)  $U(P, f) = 40$

(iv)  $L(P, f) = -40$ .

(e)  $\int_0^{\infty} \sqrt{t} e^{-t^3} dt$  is equal to

(i)  $\frac{\sqrt{\pi}}{3}$

(ii)  $\frac{\sqrt{\pi}}{2}$

(iii)  $\frac{\sqrt{\pi}}{4}$

(iv)  $2\sqrt{\pi}$ .

(f) The improper integral  $\int_1^{\infty} \frac{dx}{x^{\mu-2}}$  is convergent if and only if

(i)  $\mu = 1$

(ii)  $\mu < 2$

(iii)  $\mu \geq 2$

(iv)  $\mu > 3$ .

(g) The radius of convergence of the power series  $x + \frac{x^2}{2^2} + \frac{x^3}{3^3} + \frac{x^4}{4^4} + \dots$  is

(i)  $e$

(ii)  $\frac{1}{e}$

(iii)  $\infty$

(iv)  $0$ .

(h) The limit function of  $\left\{ \frac{x^n}{1+x^n} \right\}_n$  on  $[0, 2]$  is

(i) monotonically decreasing

(ii) monotonically increasing

(iii) continuous

(iv) not monotonic.

(i) Given that the interval of uniform convergence of a power series is  $(-4, 2)$ , for suitable  $a_n$ , which could be power series?

(i)  $\sum_{n=0}^{\infty} a_n (X+3)^n$

(ii)  $\sum_{n=0}^{\infty} a_n (X-3)^n$

(iii)  $\sum_{n=0}^{\infty} a_n (X+1)^n$

(iv)  $\sum_{n=0}^{\infty} a_n (X-1)^n$ .

(j) The sum of the Fourier series for the function  $f : [-\pi, \pi] \rightarrow \mathbb{R}$  defined by

$$f(x) = \begin{cases} -1, & -\pi \leq x < 0 \\ -2, & 0 \leq x \leq \pi \end{cases} \text{ at } x = \pi \text{ is}$$

(i)  $-\frac{1}{2}$

(ii)  $-2$

(iii)  $-\frac{3}{2}$

(iv)  $\frac{3}{2}$ .

2. Answer **any three** questions :

(a) State and prove a necessary and sufficient condition for Riemann integrability of a bounded function  $f$  defined on  $[a, b]$ . 1+4

(b) If a real-valued function  $f$  is Riemann integrable on  $[a, b]$  then prove that  $|f|$  is also Riemann integrable on  $[a, b]$  and  $\left| \int_a^b f \right| \leq \int_a^b |f|$ . 3+2

(c) (i) If  $f : [a, b] \rightarrow \mathbb{R}$  is a continuous function, such that  $f(x) \geq 0$  on  $[a, b]$  and  $\int_a^b f = 0$ , then prove that  $f$  is identically zero on  $[a, b]$ .

(ii) Prove, with justification,  $\frac{\pi^2}{9} \leq \int_{\pi/6}^{\pi/2} \frac{x}{\sin x} dx \leq \frac{2\pi^2}{9}$ . 2+3

(d) Let  $f(t) = \lim_{n \rightarrow \infty} \frac{t^n + 1}{t^n + 3}$ ,  $0 \leq t \leq 2$  and  $F(x) = \int_0^x f(t) dt$ ,  $x \geq 0$ . Prove that  $F$  is continuous at '1' but is not derivable there. 2+3

(e) (i) Prove or disprove : If  $f : [a, b] \rightarrow \mathbb{R}$  has a primitive on  $[a, b]$ , then the set of points of discontinuity of  $f$  in  $[a, b]$  is a negligible set.

(ii) Identify the set of points of discontinuity of the function  $f : [0, 1] \rightarrow \mathbb{R}$  defined by

$$f(x) = \begin{cases} \frac{1}{3^n}, & \text{when } \frac{1}{3^{n+1}} < x \leq \frac{1}{3^n} (n = 0, 1, 2, \dots) \\ 0, & \text{when } x = 0 \end{cases}$$

Hence, tell whether  $f$  is Riemann integrable on  $[0, 1]$ . 2+(2+1)

**Please Turn Over**

3. Answer **any two** questions :

(a) Let the functions  $f, g$  be positive-valued, bounded and Riemann integrable over  $[a, X]$  for every

$X > a$  such that  $\lim_{x \rightarrow \infty} \frac{f(x)}{g(x)} = \infty$ . If  $\int_a^{\infty} f$  is convergent prove that  $\int_a^{\infty} g$  is also convergent.

Is the converse true? Justify your answer. 3+2

(b) Show that  $\int_0^{\infty} \frac{\sin x}{x} dx$  is convergent but  $\int_0^{\infty} \frac{|\sin x|}{x} dx$  is not convergent. 2+3

(c) Prove that  $B(m, n) = \int_0^1 \frac{t^{m-1} + t^{n-1}}{(1+t)^{m+n}} dt$  where  $m > 0, n > 0$ . 5

(d) (i) Examine the convergence of  $\int_1^2 \frac{\log x}{\sqrt{2-x}} dx$ .

(ii) Examine the absolute convergence of  $\int_0^{\infty} \frac{\cos x dx}{\sqrt{1+x^3}}$ . 2+3

4. Answer **any four** questions :

(a) Let  $\{f_n\}_n$  be a sequence of Riemann integrable functions defined on  $[a, b]$  and  $\{f_n\}_n$  have a uniform limit  $f$  on  $[a, b]$ . Prove that  $f$  is Riemann integrable over  $[a, b]$ .

Moreover, show that  $\lim_{n \rightarrow \infty} \int_a^b f_n(x) dx = \int_a^b \left( \lim_{n \rightarrow \infty} f_n(x) \right) dx$ . 3+2

(b) Show that the sequence  $\left\{ \frac{nx}{1+n^2x^2} \right\}_n$  of continuous functions defined on  $[0, 1]$  has a continuous limit function, although the convergence is not uniform. 2+3

- (c)  $\sum_n M_n$  is a convergent infinite series of positive real numbers such that  $|f_n(x)| \leq M_n$  for all  $x \in S$  and for every  $n \in \mathbb{N}$ . Prove that  $\sum_n f_n$  is uniformly convergent on  $S$ .

Hence, prove that  $\sum_{n=1}^{\infty} \frac{\cos^3 nx}{4n^2 + 1}$  is uniformly convergent on  $[0, \infty)$ . 3+2

- (d) Examine term-by-term differentiability of  $\sum_{n=1}^{\infty} \frac{1}{n^3 + n^4 x^2}$  on  $\mathbb{R}$ . 5

- (e) Let  $\sum_{n=0}^{\infty} a_n x^n$  be a power series and  $\mu = \overline{\lim} |a_n|^{\frac{1}{n}}$ . If  $0 < \mu < \infty$ , prove that the series is absolutely convergent for  $|x| < \frac{1}{\mu}$  and is not convergent for  $|x| > \frac{1}{\mu}$ . 5

- (f) Assuming the power series for  $(1+x)^{-1}$  as  $(1+x)^{-1} = 1 - x + x^2 - x^3 + \dots$  ( $-1 < x < 1$ ), obtain the power series expansion of  $\log_e(1+x)$  and find the region of convergence of the power series of  $\log_e(1+x)$ . 3+2

- (g) Find the Fourier series of the function  $f(x) = \begin{cases} -1, & -\pi \leq x \leq 0 \\ 1, & 0 < x \leq \pi \end{cases}$ .

Also find the sum of the series at  $x = 0$  and  $x = \frac{\pi}{2}$ . 3+2

---