

(e) $(x^2 - y^2 - z^2) \frac{\partial z}{\partial x} + 2xy \frac{\partial z}{\partial y} = 2xz$ has a solution

(i) $x^2 + y^2 + z^2 = z f\left(\frac{y}{z}\right)$

(ii) $x^2 - y^2 - z^2 = y f\left(\frac{y}{z}\right)$

(iii) $x^2 + y^2 + z^2 = f\left(\frac{y}{z}\right)$

(iv) $x^2 - y^2 - z^2 = z f\left(\frac{y}{z}\right)$

(f) The complete solution of the non-linear partial differential equation $\left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2 = c^2$ is

(i) a cone

(ii) a cylinder

(iii) a sphere

(iv) none of these.

(g) Value of $\iint xy \, dx \, dy$ over the region bounded by $xy = 1, y = 0, y = x, x = 1$ is

(i) $\frac{1}{8}$

(ii) $\frac{1}{4}$

(iii) 1

(iv) $\frac{1}{2}$

(h) If the order of integration $\int_0^1 dy \int_{x=y}^{x=\sqrt{y}} f(x, y) \, dx$ is interchanged, then it will take the form

(i) $\int_0^1 dx \int_{y=x^2}^{y=x} f(x, y) \, dy$

(ii) $\int_0^1 dx \int_x^1 f(x, y) \, dy$

(iii) $\int_0^1 dx \int_{x^2}^1 f(x, y) \, dy$

(iv) none of these.

(i) If $\vec{F} = x^2 y \hat{i} + xz \hat{j} + 2yz \hat{k}$, then the value of $\text{div}\{\text{curl } \vec{F}\}$ is

(i) 1

(ii) 0

(iii) 2

(iv) $\hat{i} + \hat{k}$

(j) The work done by a particle in the force field $\vec{F} = 3x^2 \hat{i} + (2xz - y) \hat{j} + z \hat{k}$ along straight line from $(0, 0, 0)$ to $(2, 1, 3)$ is

(i) 16 units

(ii) 22 units

(iii) 14 units

(iv) 42 units.

(3)

X(4th Sm.)-Mathematics-II/CC-9/CBCS

Group – B

(Marks : 21)

Answer **any three** questions.

2. (a) Apply Charpit's method to find the complete integral of the PDE $(p + q)(px + qy) = 1$.

(b) Form a PDE by eliminating the arbitrary function ϕ and ψ from the relation $u(x, y) = y\phi(x) + x\psi(y)$.
4-3

3. Using method of separation of variables solve the PDE $4z_x + z_y = 3z$ under the condition

$$z = 3e^{-y} - e^{-5y} \text{ at } x = 0. \quad 7$$

4. Using $\eta = x + y$ as one of the transformation variable, obtain the canonical form of

$$\frac{\partial^2 u}{\partial x^2} - 2 \frac{\partial^2 u}{\partial x \partial y} + \frac{\partial^2 u}{\partial y^2} = 0$$

and hence solve it. 7

5. A tightly stretched string of length l with fixed end points is initially at rest in its equilibrium position, and each of its points is given a velocity v , which is given by

$$v(x) = \begin{cases} cx, & 0 \leq x < \frac{l}{2} \\ c(l-x), & \frac{l}{2} \leq x \leq l \end{cases}$$

Find the displacement, c being the wave speed. 7

6. Solve the following initial boundary value problem

$$u_t = u_{xx} \quad (0 < x < \lambda, t > 0)$$

subject to the conditions $u(x, 0) = 3 \sin n\pi x$ (n a+ve integer)

$$u(0, t) = u(\lambda, t) = 0. \quad 7$$

Group – C

(Marks : 24)

Answer **any four** questions.

7. Using differentiation under the sign of integration find the value of $\int_0^{\infty} e^{-a^2 x^2} \cos^2 bx \, dx$. 6

8. Evaluate the integral $\iint \frac{dx \, dy}{(1+x^2+y^2)^2}$ taken over the triangle with vertices at $(0, 0)$, $(2, 0)$ and $(1, \sqrt{3})$. 6

Please Turn Over

9. Find the value of the integral $\iiint_E \frac{dx dy dz}{x^2 + y^2 + (z-2)^2}$, where $E = \{(x, y, z) : x^2 + y^2 + z^2 \leq 1\}$. 6
10. Define conservative vector field \vec{F} and express its relation with the scalar potential $\phi(x, y, z)$. Write down Gauss's divergence theorem in case of an irrotational vector over the hemisphere centered at origin. 4+2
11. Find $\oint_c x dy + y dx$ bounded by the closed contour of astroid with $x = a \cos^3 t$ and $y = a \sin^3 t$. 6
12. Find the surface area of the region common to the intersecting cylinders $x^2 + y^2 = a^2$ and $x^2 + z^2 = a^2$. 6
13. Prove that the volume common to the sphere $x^2 + y^2 + z^2 = a^2$ and the cylinder $x^2 + y^2 = ay$ is $\frac{2}{9}(3\pi - 4)a^3$. 6
-