

2025

MATHEMATICS — HONOURS

Paper : DSCC-7

(Multivariate Calculus - I and Partial Differential Equations - I)

Full Marks : 75

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.**All symbols have their usual meanings.*

Group – A

[Multivariate Calculus – I]

(Marks : 60)

1. Answer *any five* questions :

2×5

- (a) Find the limit $\lim_{(x,y) \rightarrow (0,0)} \frac{x-y^2}{x^2+y^2}$, if it exists.
- (b) Find the equation of the tangent plane to the surface $z = \sin x + e^{xy} + 2y$ at the point $(0, 1, 3)$.
- (c) Verify whether the set $S = \{(x, y) \in \mathbb{R}^2 : 0 < x < 1, 0 < y < 1\}$ is an open set in \mathbb{R}^2 or not.
- (d) Evaluate $\iint_S (x^2 + y^2) dS$, where $S : z = 4 - x - 2y, 0 \leq x \leq 4, 0 \leq y \leq 2$.
- (e) If $x = u^2v^2, y = v^2 + u^2$, find the Jacobian $\frac{\partial(x,y)}{\partial(u,v)}$.
- (f) If $f = x^3 + y^3 + z^3$, find the directional derivative of f at the point $(1, -1, 2)$ in the direction of the vector $\hat{j} - \hat{k}$.
- (g) If $v = f(u)$, where u is a homogeneous function of two independent variables x, y of degree n , then prove that $x \frac{\partial v}{\partial x} + y \frac{\partial v}{\partial y} = nu \frac{dv}{du}$.
- (h) If $f(t) = \int_0^t \sqrt{1+s^2} ds, s > 0$, then find $f'(2)$.

Please Turn Over

(2886)

2. Answer **any five** questions :

(a) (i) Let (x, y) approach towards $(0, 0)$ along the line $y = -x$. Using Taylor's theorem for function

of two variables, show that $\lim_{\substack{(x,y) \rightarrow (0,0) \\ (y=-x)}} \frac{\sin xy + xe^x - y}{x \cos y + \sin 2y} = -2$.

(ii) Transform the equation $x^2 \frac{\partial z}{\partial x} + y^2 \frac{\partial z}{\partial y} = z^2$, by introducing new independent variable

$u = x, v = \frac{1}{y} - \frac{1}{x}$ and the new function $w = \frac{1}{z} + \frac{1}{x}$. 5+5

(b) (i) Let $f(x, y) = \begin{cases} \frac{xy^3}{x^2 + y^2}, & (x, y) \neq (0, 0) \\ 0, & (x, y) = (0, 0) \end{cases}$. Show that $f_{yx}(0, 0) \neq f_{xy}(0, 0)$. Which condition of

Schwarz's theorem is not satisfied for this f ?

(ii) State and prove Euler's theorem on homogeneous function of three variables.

(4+1)+(1+4)

(c) (i) Use transformation $x = \frac{u}{v}, y = v$ to evaluate the integral $\iint_R xy \, dA$, where R is the region in the first quadrant bounded by the lines $y = x$ and $y = 3x$ and the hyperbolas $xy = 1, xy = 3$.

(ii) Evaluate $\iiint_E (9 - x^2 - y^2) dV$, where E is the solid hemisphere $x^2 + y^2 + z^2 \leq 9, z \geq 0$.

5+5

(d) (i) Find the area of the surface $z = \frac{2}{3} \left(x^{\frac{3}{2}} + y^{\frac{3}{2}} \right)$, in the region $0 \leq x \leq 1, 0 \leq y \leq 1$.

(ii) By changing the order of integration, prove that

$$\int_0^1 dx \int_x^{\frac{1}{x}} \frac{y \, dy}{(1+xy)^2(1+y^2)} = \frac{(\pi-1)}{4}.$$

5+5

(3)

D(4th Sm.)-Mathematics-H/DSCC-7/CCF

- (e) (i) If $u = \tan^{-1} \frac{x^3 + y^3}{x - y}$, then prove that $x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = \sin 2u (1 - 4 \sin^2 u)$.
- (ii) Investigate the maxima and minima of the function $f(x, y) = x^2 + 4xy + 4y^2 + x^3 + 2x^2y + y^4$.
5+5
- (f) (i) Assuming the validity of differentiation under the sign of integration, show that

$$\int_0^{\frac{\pi}{2}} \frac{\log(1 + y \sin^2 x)}{\sin^2 x} dx = \pi(\sqrt{1+y} - 1), \text{ where } y > -1.$$

- (ii) Evaluate $\iint \frac{dx dy}{(1 + x^2 + y^2)^2}$ over a loop of the lemniscate $(x^2 + y^2)^2 - (x^2 - y^2) = 0$. 5+5
- (g) (i) Prove Taylor's theorem for a function $f(x, y)$ of two variables stating the conditions imposed upon $f(x, y)$.
- (ii) Using the method of Lagrange's multiplier show that the minimum value of $bcx + cay + abz$ subject to $xyz = abc$ is $3abc$. 5+5
- (h) (i) Let $f: D \rightarrow \mathbb{R}$ be a function where $D \subset \mathbb{R}^2$. When is a function $f(x, y)$ said to be differentiable at the point (a, b) in $D \subset \mathbb{R}^2$? Show that differentiability of f at (a, b) implies the continuity of f at (a, b) and existence of both f_x and f_y at that point.
- (ii) What do you mean by an implicit function $y = \phi(x)$ defined by $F(x, y) = 0$ near (a, b) ? Verify implicit function theorem for $x^2 + xy + y^2 - 1 = 0$ near $(0, -1)$. (1+2+2)+(2+3)

Group - B

[Partial Differential Equations - I]

(Marks : 15)

3. Answer **any three** questions :

- (a) Solve the partial differential equation $(x^2 - yz) \frac{\partial z}{\partial x} + (y^2 - zx) \frac{\partial z}{\partial y} = z^2 - xy$ by Lagrange's method. 5
- (b) Find the PDE arising from the family of surfaces $z = xy + f(x^2 + y^2)$, f being an arbitrary function. 5
- (c) Find the complete integral of $p^2x + q^2y = z$ by Charpit's method, where $p \equiv \frac{\partial z}{\partial x}$, $q \equiv \frac{\partial z}{\partial y}$. 5

Please Turn Over

(2886)

- (d) Find the integral surface of the quasi linear PDE $yp - 2xyq = 2xz$ which passes through the curve $x = t, y = t^2, z = t^3$, where $p = \frac{\partial z}{\partial x}, q = \frac{\partial z}{\partial y}$. 5
- (e) Obtain the solution of the Cauchy problem $xu_x + yu_y = u + 1$ with $u(x, y) = x^2$ on $y = x^2$. 5
-